

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 1 209 151 A1

(12)

EUROPEAN PATENT APPLICATION

published in accordance with Art. 158(3) EPC

(43) Date of publication:

29.05.2002 Bulletin 2002/22

(21) Application number: 00956885.8

(22) Date of filing: 01.09.2000

(51) Int Cl⁷: C07D 211/32, C07D 401/06,
C07D 498/04, C07D 417/06,
C07D 413/06, A61K 31/445,
A61K 31/55, A61P 25/28,
A61K 31/454, A61P 25/14,
A61P 43/00

(86) International application number:

PCT/JP00/05968

(87) International publication number:

WO 01/16105 (08.03.2001 Gazette 2001/10)

(84) Designated Contracting States:

AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE

(30) Priority: 01.09.1999 JP 24711599

(71) Applicant: Eisai Co., Ltd.

Tokyo 112-8089 (JP)

(72) Inventors:


- IIMURA, Yoichi
Tsukuba-shi, Ibaraki 305-0051 (JP)
- KOSASA, Takashi
Ibaraki 300-2436 (JP)

(74) Representative: HOFFMANN - EITLE

Patent- und Rechtsanwälte
Arabellastrasse 4
81925 München (DE)

(54) 4-SUBSTITUTED PIPERIDINE DERIVATIVES

(57) The present invention provides a novel compound having an excellent acetylcholinesterase inhibitory effect. That is, it provides a compound represented by formula:



(I)

(wherein R¹ and R² represent substituents), a pharmacologically acceptable salt thereof or hydrates thereof.

EP 1 209 151 A1

Description

Field of the Invention

[0001] The present invention relates to a novel compound useful as an acetylcholinesterase inhibitor, a salt thereof and hydrates thereof, and to a process for producing them.


Prior Art

[0002] It is known that senile dementia such as Alzheimer-type senile dementia, cerebrovascular dementia, attention deficit hyperactivity disorder etc. are accompanied with cholinergic hypofunction. Currently, acetylcholinesterase inhibitors have been acknowledged to be useful as a remedy for these diseases and actually used in clinical applications. For example, as their typical remedies, there have been known donepezil hydrochloride (1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine hydrochloride), and also rivastigmine (3-[1-(dimethylamino)ethyl]phenyl N-ethyl-N-methylcarbamate), metrifonate (dimethyl (2,2,2-trichloro-1-hydroxyethyl)phosphate), tacrine hydrochloride (1,2,3,4-tetrahydro-9-acridinamine), galanthamine hydrobromide, neostigmine, physostigmine etc.


[0003] However, among the aforementioned medicaments, it is only donepezil hydrochloride that has been acknowledged sufficient usefulness in consideration of pharmacological activity, side effect, number of administrations, form of administration etc., and short of it no acetylcholinesterase inhibitor that is clinically useful has been found yet. Although donepezil hydrochloride is an excellent medicine, it has been keenly desired to provide besides donepezil hydrochloride an acetylcholinesterase inhibitor that has excellent effect and is highly useful since an acetylcholinesterase inhibitor that exhibits more excellent effect, if available, will offer a wider choice of options in selecting medicines in clinical fields.










Disclosure of the Invention

[0004] Under these circumstances, the present inventors have made intensive research over the years in order to develop medicines having more excellent effects and higher safety. As a result, they have succeeded in synthesizing a novel compound represented by the formula:




wherein, R¹ represents any group selected from the groups represented by formulae:






and



(wherein (R³)s are the same as or different from each other and each represents hydrogen atom, a halogen atom, hydroxyl group, a C₁₋₆ alkyl group, a C₂₋₈ cycloalkyl group, a C₁₋₆ alkoxy group, a C₁₋₆ alkoxyalkoxy group, a halogeno C₁₋₆ alkyl group, a hydroxy C₁₋₆ alkyl group, a cyano C₁₋₆ alkyl group, an amino C₁₋₆ alkyl group, a halogeno C₁₋₆ alkoxy group, a hydroxy C₁₋₆ alkoxy group, a cyano C₁₋₆ alkoxy group, a lower acyl group, nitro group, an optionally substituted amino group, an optionally substituted carbamoyl group, mercapto group or a C₁₋₆ thioalkoxy group; R⁴ represents hydrogen atom or a C₁₋₆ alkyl group; R⁵ represents a halogen atom (provided that fluorine is excluded), hydroxy group, a C₁₋₆ alkyl group, a C₁₋₆ alkoxy group, cyano group, a halogeno C₁₋₆ alkyl group, a hydroxy C₁₋₆ alkyl group, a cyano C₁₋₆ alkyl group, an amino C₁₋₆ alkyl group, nitro group, an optionally substituted amino group, an optionally substituted carbamoyl group, an optionally substituted carboxyl group, mercapto group or a C₁₋₆ thioalkoxy group; the partial structure --- represents a single bond or double bond; m is 0 or an integer from 1 to 6; n is an integer from 1 to 4; and p is an integer of 1 or 2); and

R² represents a C₃₋₈ cycloalkylmethyl, a 2,2-(alkylenedioxy)ethyl or a group represented by the formula:




wherein the ring A represents a benzene ring or a heterocyclic ring; (R⁶)s are the same as or different from each other and each represents hydrogen, a halogen atom, hydroxyl group, nitrile group, a C₁₋₆ alkyl group, a C₃₋₈ cycloalkyl group, a C₁₋₆ alkoxy group, a C₁₋₆ alkoxyalkoxy group, an aryloxy group, an aralkyloxy group, a halogeno C₁₋₆ alkyl group, a hydroxy C₁₋₆ alkyl group, a cyano C₁₋₆ alkyl group, a halogeno C₁₋₆ alkoxy group, a hydroxy C₁₋₆ alkoxy group, a cyano C₁₋₆ alkoxy group, a lower acyl group, nitro group, an optionally substituted amino group, an optionally substituted amide group, mercapto group or a C₁₋₆ thioalkoxy group, and two of the R⁶ may together form an aliphatic ring, an aromatic ring, a heterocyclic ring or an alkylenedioxy ring, and q is 0 or an integer from 1 to 5. Further, they have found that the compound represented by the above formula (1), a salt thereof and hydrates thereof exhibit an excellent acetylcholinesterase inhibitory effect and the intended objects have been attained therewith. Thus, they have achieved the present invention.

[0005] That is, the first feature of the present invention is:

(1) the compound represented by the above formula (1), a pharmacologically acceptable salt thereof or hydrates thereof, and further;

(2) in the above (1), R¹ may be a group represented by the formula:




wherein R^{3a} and R^{3b} are the same as or different from each other and each represents a C₁₋₆ alkoxy group; and R⁵ and m have the same meanings as defined above;

(3) in the above (2), R^{3a} and R^{3b} may be methoxy group;

(4) in the above (1) to (3), R⁵ may be chlorine or bromine;

(5) in the above (1) to (4), m may be 1;

(6) in the above (1), R² may be a group represented by the formula:



wherein the ring A, R⁶ and q have the same meanings as defined above;

(7) in the above (6), the ring A may be a benzene ring;

(8) in the above (6), the ring A may be a pyridine ring;

(9) in the above (6), q may be an integer of 1 or 2; and

(10) in the above (1), the compound may be any compound selected from:

- 1-benzyl-4-[(5,6-dimethoxy-2-chloro-1-indanon)-1-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-iodo-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-hydroxy-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-methyl-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-ethyl-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-azido-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-amino-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-methylamino-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-dimethylamino-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-acetamide-1-indanon)-2-yl]methylpiperidine,
- 1-benzyl-4-[(5,6-dimethoxy-2-methanesulfonamide-1-indanon)-2-yl]methylpiperidine,
- 3-(1-benzylpiperidin-4-yl)-2-chloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone,
- 3-(1-benzylpiperidin-4-yl)-2-dichloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone,
- 5,7-dihydro-3-(1-chloro-2-[(1-(phenylmethyl)-4-piperidinyl)ethyl]-6H-pyrrolo[4,5-f]-1,2-benzisocoxazol-

6-one,
 5,7-dihydro-3-[1,1-dichloro-2-[(1-(phenylmethyl)-4-piperidinyl)ethyl]-6H-pyrrolo[4,5-f]-1,2-benzooxazol-
 6-one,
 1-(2-methyl-6-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2-chloro-1-propanone, and
 1-(2-methyl-6-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2,2-dichloro-1-propanone.

[0006] The second feature of the present invention is:

(11) a medicament containing the compound described in the above (1), a salt thereof or hydrates thereof, and further,

(12) the medicament in the above (11) may be an acetylcholinesterase inhibitor;

(13) the medicament in the above (11) may be an agent for treating, preventing or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder; and

(14) the senile dementia in the above (13) may be Alzheimer-type senile dementia.

[0007] The present invention provides a medical composition comprising a pharmacologically effective amount of the compound described in the above (1), a pharmacologically acceptable salt thereof or hydrates thereof, and a pharmacologically acceptable carrier.

[0008] Also, the present invention provides a method for preventing, treating or ameliorating diseases against which the inhibition of an acetylcholinesterase is effective and a method for treating, preventing or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder, by administering a pharmacologically effective amount of the compound described in the above (1), a pharmacologically acceptable salt thereof or hydrates thereof to a patient.

[0009] Further, the present invention provides use of the compound described in the above (1), a pharmacologically acceptable salt thereof or hydrates thereof for producing an agent for preventing, treating or ameliorating diseases against which the inhibition of an acetylcholinesterase is effective, for producing an acetylcholinesterase inhibitor, and for producing an agent for preventing, treating or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder.

[0010] Hereinafter, the symbols, terms, etc., used in the specification of the present application will be explained and the present invention will be illustrated in detail.

[0011] Note that the structural formula for a compound as used in the specification of the present application may be expressed as indicating a certain isomer or isotope for the sake of convenience. However, in the present invention, a compound represented by a structural formula is to be construed as including all the possible isomers such as geometric isomers, optical isomers due to asymmetric carbon atom, stereoisomers and tautomers and mixtures thereof as well as isotopes and should not be limited to what is expressed by the formula that is presented only for the convenience sake. The compound may be any one of isomers or mixtures thereof, or any one of isotopes. Therefore, a compound may have an asymmetric carbon atom or atoms in the molecule and hence there may be optically active forms or racemic form. In the present invention compounds are not limited to particular ones but include both of them. Further, crystal polymorphism may exist and similarly the present invention is not limited to a particular crystal form but may include any one of crystal forms or mixtures thereof. The compounds of the present invention may be hydrates as well as anhydrides.

[0012] In the formula (I) above, the "halogen atom" represented by R³ in the definition of R¹ indicates an atom such as fluorine, chlorine, bromine or iodine, preferably fluorine, chlorine or bromine. The "halogen atom" represented by R⁵ includes halogen atoms other than fluorine and includes, for example, chlorine, bromine, iodine and the like, and preferably chlorine and bromine.

[0013] The "C₁₋₆ alkyl group" represented by R³, R⁴ and R⁵ above means an alkyl group having 1 to 6 carbon atoms and includes linear or branched alkyl groups, for example, methyl group, ethyl group, n-propyl group, i-propyl group, n-butyl group, i-butyl group, t-butyl group, n-pentyl group, i-pentyl group, neopentyl group, hexyl group, 1-methylpropyl group, 1-methylbutyl group, 2-methylbutyl group etc.

[0014] The "C₃₋₆ cycloalkyl group" represented by R³ above means a cyclic alkyl group having 3 to 6 carbon atoms and includes, for example, a cyclopropyl group, a cyclobutyl group, a cyclopentyl group, a cyclohexyl group, a cycloheptyl group, a cyclooctyl group etc.

[0015] The "C₁₋₆ alkoxy group" represented by R³ and R⁵ above means a group consisting of a group having the same meaning as the "C₁₋₆ alkyl group" defined above and an oxygen atom bonded thereto and includes linear or branched alkoxy groups, for example, methoxy group, ethoxy group, n-propoxy group, i-propoxy group, n-butoxy group, i-butoxy group, pentyloxy group, hexyloxy group etc.

[0016] The "C₁₋₆ alkoxyalkoxy group" represented by R³ above means a group consisting of a group having the same meaning as the "C₁₋₆ alkoxy group" defined above and another "C₁₋₆ alkoxy group" bonded thereto and includes,

for example, methoxy-methoxy group, methoxyethoxy group, methoxypropoxy group, ethoxymethoxy group, ethoxyethoxy group, ethoxypropoxy group, propoxypropoxy group etc.

[0017] The "halogeno C_{1-6} alkyl group" represented by R^3 and R^5 above means a group consisting of a group having the same meaning as the C_{1-6} alkyl group defined above having one or more halogen atoms bonded thereto which may be the same as or different from and includes, for example, chloromethyl group, dichloromethyl group, trichloromethyl group, fluoromethyl group, difluoromethyl group, trifluoromethyl group, fluoromethyl group, difluoromethyl group, trifluoromethyl group etc.

[0018] The "hydroxy C_{1-6} alkyl group" represented by R^3 and R^5 above means a group consisting of a group having the same meaning as the C_{1-6} alkyl group defined above having one or more hydroxyl groups bonded thereto and includes, for example, hydroxymethyl group, hydroxyethyl group, 2,3-dihydroxypropyl group etc.

[0019] The "cyano C_{1-6} alkyl group" represented by R^3 and R^5 above means a group consisting of a group having the same meaning as the C_{1-6} alkyl group defined above having bonded thereto one or more cyano groups bonded thereto and includes, for example, cyanomethyl group, cyanoethyl group, cyanoethyl group, cyanoethyl group etc.

[0020] The "halogeno C_{1-6} alkoxy group" represented by R^3 above means the "halogeno C_{1-6} alkyl group" having the same meaning as the halogeno C_{1-6} alkyl group defined above, which is bonded to an oxygen atom, the "hydroxy C_{1-6} alkoxy group" means the "hydroxy C_{1-6} alkyl group" having the same meaning as the hydroxy C_{1-6} alkyl group as defined above, which is bonded to an oxygen atom, and the "cyano C_{1-6} alkoxy group" means the "cyano C_{1-6} alkyl group" having the same meaning as the cyano C_{1-6} alkyl group as defined above, which is bonded to an oxygen atom.

[0021] The "lower acyl group" represented by R^3 above means a linear or branched acyl group derived from a fatty acid having 1 to 6 carbon atoms and includes, for example, formyl group, acetyl group, propionyl group, butyryl group, isobutyryl group, valeryl group, isovaleryl group, pivaloyl group, hexanoyl group etc.

[0022] The "optionally substituted amino group" represented by R^3 and R^5 above means an amino group of which the nitrogen atom may be substituted by a group such as a C_{1-6} alkyl group and further the amino group includes cyclic amino groups. The "optionally substituted amino group" includes, for example, amino group ($-NH_2$), methylamino group ($-NHCH_3$), dimethylamino group ($-N(CH_3)_2$), pyrrolidinyl group, pyrazolinyl group, piperidinyl group, piperazinyl group etc.


[0023] The "optionally substituted carbamoyl group" represented by R^3 and R^5 above means an amide group of which the nitrogen atom may be substituted by a group such as a C_{1-6} alkyl group and further the amide group includes amide groups of cyclic amine. The "optionally substituted amide group" includes, for example, amide group ($-CONH_2$), N-methylamide group ($-CONHCH_3$), N,N-dimethylamide group ($-CON(CH_3)_2$), N-ethylamide group ($-CONHC_2H_5$), N,N-diethylamide group ($-CON(C_2H_5)_2$), N-methyl-N-ethylamide group ($-CON(CH_3)(C_2H_5)$), pyrrolidinylcarbonyl group, pyrazolinylcarbonyl group, piperidinylcarbonyl group, piperazinylcarbonyl group etc.

[0024] The " C_{1-6} thioalkoxy group" represented by R^3 and R^5 above means a group consisting of a group having the same meaning as the C_{1-6} alkyl group defined above bonded to a sulfur atom and includes, for example, methylthio group ($-SCH_3$), ethylthio group ($-SC_2H_5$) etc.


[0025] In the formula (I) above, the symbol m in R^1 indicates 0 or an integer from 1 to 6, m is preferably 0 or an integer from 1 to 5, more preferably 0 or an integer from 1 to 3, still more preferably 0 or an integer of 1 or 2 and most preferably 0 or

1. The symbol n indicates an integer of 1 to 4, n is preferably an integer from 1 to 3 and more preferably an integer of 1 or
2. Further, the symbol p indicates an integer of 1 or 2, p is preferably 1.

[0026] In the above formula (I), preferred groups represented by R^1 are those groups represented by the formula:



wherein the symbols have the same meanings as defined above. More preferably, mention may be made of those groups represented by the formula:



wherein R^{3a} , R^{3b} , R^5 and m have the same meanings as defined above. In this case, further preferred groups include those groups in which R^{3a} and R^{3b} represent methoxy groups. The most preferred groups in R^1 include a [(5,6-dimethoxy-2-chloro-1-indanon)-2-yl]methyl group, a [(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methyl group and the like.

[0027] The " C_{3-6} cycloalkyl group" in the " C_{3-6} cycloalkylmethyl group" represented by R^2 in the above formula (I) means a group having the same meaning as the C_{3-6} cycloalkyl group defined above and the " C_{3-6} cycloalkylmethyl group" means a group consisting of the above-mentioned " C_{3-6} cycloalkyl group" and a methyl group bonded thereto, which includes, for example, cyclopropylmethyl group, cyclobutylmethyl group, cyclopentylmethyl group, cyclohexylmethyl group, cycloheptylmethyl group, cyclooctylmethyl group and the like. Among these, cyclohexylmethyl group is more preferred.

[0028] The " $2,2-(alkylenedioxy)ethyl$ group" represented by R^2 above means a group (acetal group) consisting of ethyl group, the terminal carbon atom of which is substituted with a cyclic alkylenedioxy group and includes, for example, 2,2-(ethylenedioxy)ethyl group (also known as {1,3-dioxolan-2-yl}methyl group), 2,2-(propylenedioxy)ethyl group (also known as {1,3-dioxane-2-yl}methyl group), 2,2-(butylenedioxy)ethyl group (also known as {1,3-dioxepan-2-yl}methyl group) etc. Among these, 2,2-(ethylenedioxy)ethyl group is preferred.

[0029] The "aryl group" in the "aryloxy group" represented by R^6 above means a cyclic hydrocarbon group that constitutes an aromatic group and includes monocyclic, bicyclic or tricyclic aryl groups such as phenyl group, indenyl group, naphthyl group, azulenyl group, heptalenyl group, anthranyl group and phenanthrenyl group. The "aryloxy group" means a group consisting of the above-mentioned aryl group and oxygen atom bonded thereto and includes, for example, phenoxy group, naphthyloxy group etc.

[0030] The "aralkyloxy group" represented by R^6 above means a group consisting of a group having the same meaning as the aryl group and a C_{1-6} alkyl group bonded thereto and the resulting aralkyl group being further bonded to oxygen atom and includes, for example, benzyloxy group, phenylethoxy group, phenylpropoxy group, naphthylmethoxy group etc.


[0031] The "halogen atom", " C_{1-6} alkyl group", " C_{3-6} cycloalkyl group", " C_{1-6} alkoxy group", " C_{1-6} alkoxyalkoxy group", "halogeno C_{1-6} alkyl group", "hydroxy C_{1-6} alkyl group", "cyano C_{1-6} alkyl group", "halogeno C_{1-6} alkoxy group", "hydroxy C_{1-6} alkoxy group", "cyano C_{1-6} alkoxy group", "lower acyl group", "optionally substituted amino group", "optionally substituted amide group" and " C_{1-6} thioalkoxy group" have the same meanings as the halogen atom, C_{1-6} alkyl group, C_{3-6} cycloalkyl group, C_{1-6} alkoxy group, C_{1-6} alkoxyalkoxy group, halogeno C_{1-6} alkyl group, hydroxy C_{1-6} alkyl group, cyano C_{1-6} alkyl group, halogeno C_{1-6} alkoxy group, hydroxy C_{1-6} alkoxy group, cyano C_{1-6} alkoxy group, lower acyl group, optionally substituted amino group, optionally substituted amide group and C_{1-6} thioalkoxy group as defined above, respectively. The C_{1-6} alkoxy carbonyl group means a group consisting of the above-mentioned C_{1-6} alkoxy group and carbonyl group bonded thereto and specifically includes, for example, methoxycarbonyl group ($-COOCH_3$), ethoxycarbonyl group ($-COOC_2H_5$) etc.

[0032] In the definition of R^3 above, examples of the fatty acid formed by two R^6 include cyclopentane ring, cyclohexane ring, cycloheptane ring, cyclooctane ring etc. Further, examples of the aromatic ring formed by two R^6 include benzene ring. Moreover, examples of the heterocyclic ring formed by two R^6 include furan ring, thiophene ring, pyrrole ring, imidazole ring, oxazole ring, thiazole ring, triazole ring, pyridine ring, pyrazine ring, pyrimidine ring, tetrahydropyran ring, tetrahydropyran ring, dioxane ring, dioxolane ring, piperidine ring, piperazine ring, morpholine ring, thiomorpholine ring etc. Examples of the alkylenedioxy ring formed by two R^6 include methylenedioxy group, ethylenedioxy group, propylenedioxy group etc.


[0033] The "heterocyclic ring" represented by the ring A in R^2 above means a ring containing 1 to 4 hetero atoms such as nitrogen atom, sulfur atom or oxygen atom and includes a "5- to 14-membered aromatic heterocyclic ring" and a "5- to 10-membered non-aromatic heterocyclic ring". The "5- to 14-membered aromatic heterocyclic ring" includes monocyclic, bicyclic or tricyclic 5- to 14-membered aromatic heterocyclic rings containing 1 to 4 atoms selected from nitrogen atom, sulfur atom and oxygen atom, for example, (1) nitrogen-containing aromatic heterocyclic rings such as pyrrole, pyridine, pyridazine, pyrimidine, pyrazine, pyrazole, imidazole, indole, isindole, indolizine, purine, indazole, quinoline, isoquinoline, quinaldine, phthalazine, naphthyridine, quinoxaline, quinoxaline, cinnoline, pteridine, imidazo-triazine, pyrazinopyridazine, acridine, phenanthridine, carbazole, carbazoline, perimidine, phenanthroline and phenazine rings; (2) sulfur-containing aromatic heterocyclic rings such as thiophene and benzothioephene rings; (3) oxygen-containing aromatic heterocyclic rings such as furan, pyran, cyclopentapyran, benzofuran and isobenzofuran rings;

(4) aromatic heterocyclic rings containing 2 or more hetero atoms selected from nitrogen atom, sulfur atom and oxygen atom, such as thiazole, isothiazole, benzothiazole, benzothiadiazole, phenothiazine, isoxazole, furazane, phenoxazine, pyrazolooxazole, imidazothiazole, thienoturan, furanopyrrole and pyridoxazine rings. The "5- to 10-membered non-aromatic heterocyclic ring" means a hydrocarbon ring of which 1 to 4 carbon atoms are substituted with any one of hetero atoms selected from nitrogen atom, sulfur atom and oxygen atom, it being intended to include also an unsaturated condensed ring. The "5- to 10-membered non-aromatic heterocyclic ring" includes, for example, pyrrolidine, pyrroline, piperidine, piperazine, imidazoline, pyrazolidine, imidazolidine, morpholine, tetrahydropyran, aziridine, oxirane and oxathiolane rings, phthalimide, succinimide and the like. Preferred rings in the ring A include benzene, pyridine, pyridazine, pyrimidine, pyrazine, piperidine, piperazine and morpholine rings.

[0034] in the above formula (I), a preferred example in R^6 includes a group represented by the formula:



(wherein ring A, R^6 and q have the same meanings as defined above), and a more preferred example includes a group represented by the formula:



(wherein R^6 and q have the same meanings as defined above).

[0035] According to preferred modes of the present invention, the compound represented by the above formula (I) includes any compound selected from, for example, 1-benzyl-4-[(5,6-dimethoxy-2-chloro-1-indanon)-1-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-iodo-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-hydroxy-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-methyl-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-ethyl-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-azido-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-amino-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-methylamino-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-dimethylamino-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-acetamido-1-indanon)-2-yl]methylpiperidine, 1-benzyl-4-[(5,6-dimethoxy-2-methanesulfonamido-1-indanon)-2-yl]methylpiperidine, 3-(1-benzylpiperidin-4-yl)-2-chloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone, 3-(1-benzylpiperidin-4-yl)-2,2-dichloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone, 5,7-dihydro-3-(1-chloro-2-[(1-phenylmethyl)-4-piperidinyl]ethyl)-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one, 5,7-dihydro-3-(1,1-dichloro-2-[(1-phenylmethyl)-4-piperidinyl]ethyl)-6H-pyrrolo[4,5-f]-1,2-benzisoxazol-6-one, 1-(2-methyl-6-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2,2-dichloro-1-propanone, 1-(2-methyl-6-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2,2-dichloro-1-propanone and the like. Of course, the present invention is not limited to the aforementioned modes.

[0036] The "pharmacologically acceptable salt" as referred to in the present invention is not particularly limited as far as it forms an addition salt together with the compound of the present invention. For example, hydrohalogenic acid salts such as hydrofluoride, hydrochloride, hydrobromide and hydroiodide; inorganic acid salts such as sulfate, nitrate, perchlorate, phosphate, carbonate and bicarbonate; organic carboxylic acid salts, such as acetate, oxalate, maleate, tartarate and fumarate; organic sulfonic acid salts, such as methanesulfonate, trifluoromethanesulfonate, ethanesulfonate, benzenesulfonate, toluenesulfonate and camphorsulfonate; amino acid salts such as aspartate and glutamate; salts with amines such as trimethylamine salts, triethylamine salts, procaine salts, pyridine salts and phenethylbenzylamine salts; alkali metal salts such as sodium salts and potassium salts; alkaline earth metal salts such as magnesium salts and calcium salts, and hydrochloride and oxalate are preferred.

[0037] In the production of the compounds represented by the above formula (I) according to the present invention, the starting compound in the reaction may be in the form of a salt or hydrate, and is not particularly limited as far as it does not inhibit the reaction. In the case where the compounds (I) of the present invention are obtained as free forms, they can be converted into the state of salts that the compounds (I) of the present invention may form according to a conventional method. Various isomers obtained on the compound (I) of the present invention may be purified and

isolated by use of common separation means (for example, recrystallization, chromatography, etc.). In particular, in the case where optically active forms of the compounds of the present invention are needed, they can be obtained, for example, by use of a method using an optically active reagent (asymmetric synthesis), a method performing optical resolution of racemic form, or the like methods.

[0038] The compounds represented by the above formula (I) according to the present invention can be formulated into agents such as tablets, powder, fine granules, granules, coated tablets, capsules, syrups, troches, inhalants, suppositories, ointments, ophthalmic ointments, ophthalmic solutions, nose drops, ear drops, poultices, lotions etc. by a common method. In the formulation, use can be made of usually used fillers, binders, lubricants, colorants, flavoring agents, and if needed, also stabilizers, emulsifiers, absorbent agents, surfactants, pH adjusting agents, preservatives, antioxidants etc. Components generally used as starting materials for medical preparations are blended and formulated by a conventional method. For example, for producing oral preparations, the compound of the present invention or a pharmacologically acceptable salt thereof and fillers, and if necessary a binder, a disintegrant, a lubricant, a colorant, a flavoring agent etc. are added and then formulated into powder, fine granules, granules, tablets, coated tablets, capsules etc. by a conventional method. Such components include, for example, animal and plant oils such as soybean oil, beef tallow and synthetic glyceride; hydrocarbons such as liquid paraffin, squalene and solid paraffin; ester oils such as octyl decadecyl myristate and isopropyl myristate; higher alcohol such as cetostearyl alcohol and behenyl alcohol; silicone resin; silicone oil; surfactants such as polyoxyethylene fatty acid esters, sorbitan fatty acid esters, glycerol fatty acid esters, polyoxyethylene sorbitan fatty acid esters, polyoxyethylene hardened castor oil and polyoxyethylene polyoxypropylene block copolymers; water-soluble polymers such as hydroxyethylcellulose, polyacrylic acid, carboxyvinyl polymer, polyethylene glycol, polyvinylpyrrolidone and methylcellulose; lower alcohols such as ethanol and isopropanol; polyhydric alcohols such as glycerol, propylene glycol, dipropylene glycol and sorbitol; sugars such as glucose and sucrose; inorganic powder such as silicic anhydride, aluminummagnesium silicate and aluminum silicate; purified water and the like. The fillers which can be used includes, for example, lactose, corn starch, white sugar, glucose, mannitol, sorbit, crystalline cellulose, silicon dioxide etc. The binder which can be used includes, for example, polyvinyl alcohol, polyvinyl ether, methylcellulose, ethylcellulose, gum arabic, tragacanth, gelatin, shellac, hydroxypropylmethylcellulose, hydroxypropylcellulose, polyvinylpyrrolidone, polypropylene glycol/polyoxyethylene block polymer, meglumine etc. The disintegrant which can be used includes, for example, starch, agar, gelatin powder, crystalline cellulose, calcium carbonate, sodium hydrogen carbonate, calcium citrate, dextrin, pectin, carboxymethylcellulose calcium and the like. The lubricant which can be used includes, for example, magnesium stearate, talc, polyethylene glycol, silica, hardened plant oil and the like. The colorant which can be used includes those of which addition to pharmaceutical preparations is permitted. The corrigent which can be used includes cocoa powder, peppermint camphor, empass, peppermint oil, Borneo camphor, cinnamon powder and the like. The tables and granules may of course have a glycoalyx or like coating as needed. When liquid preparations such as syrup and a preparation for injection are produced, a pH adjustor, a dissolving agent, an isotonic agent and so on and optionally a dissolution auxiliary, a stabilizer and the like as needed are added to the compound of the present invention or its pharmacologically acceptable salt, and the mixture is formulated into a preparation by a conventional method. The process for producing external preparations is not limited and a conventional method may be used for producing them. That is, the base material used upon making preparations may include various raw materials that can be usually used for drugs, quasi-drugs, cosmetics and the like. The base material which can be used specifically includes, for example, animal and plant oils, mineral oils, ester oils, waxes, higher alcohols, fatty acids, silicone oil, surfactants, phospholipids, alcohols, polyhydric alcohols, water-soluble polymers, clay minerals, purified water and the like raw materials. Further, as necessary, pH adjusting agents, antioxidant, chelating agents, antiseptical fungicidal agents, coloring agents, flavoring agents and the like may be added thereto. However, the base materials for external preparations of the present invention are not limited to these materials. Also, components having activity of differentiation, bloodstream promoting agents, germicides, antiphlogistics, cell activators, vitamins, amino acids, moisturizers, keratin solubilizers and the like components may be blended as needed. The addition amounts of the base materials are usually those amounts that will give predetermined concentrations of the components upon production of external preparations.


[0039] When administering an agent containing the compound represented by the above formula (I) according to the present invention, a salt thereof or hydrates thereof as an active ingredient, the form of the agent is not particularly limited and it may be either orally or parenterally administered by a method usually used. For example, the agent may be formulated and administered as tablets, powder, granules, capsules, syrups, troches, inhalant, suppository, injection, ointment, ophthalmic ointments, eye drops, nose drops, ear drops, poultice, lotions etc. The dosage of the pharmaceutical preparation of the present invention may be selected as appropriate depending on the severity of symptom, the age, sex, body weight, the route of the administration, the type of salt, specific kinds of the disease and so on.

[0040] The following Reference Examples, Examples (further their pharmacologically acceptable salt thereof, hydrates thereof, and a medicament or medical compositions containing them), test examples are given for illustration purposes and the compounds of the present invention should by no means be construed as being limited to the following specific examples.

Reference Example 1

1-Benzyl-4-[(5-ethoxy-6-methoxy-1-indanon)-2-yl]methylpiperidine

[0041]



[0042] In 20 ml of THF was dissolved 0.20 g (0.55 mmol) of 1-benzyl-4-[(5-hydroxy-6-methoxy-1-indanon)-2-yl]methylpiperidine, followed by adding 0.064 ml (1.1 mmol) of ethanol, 0.29 g (1.1 mmol) of triphenylphosphine and 0.17 ml (1.1 mmol) of diethyl azodicarboxylate. After stirring at room temperature overnight, the mixture was evaporated. To the resulting residue was added 30 ml of water, followed by extracting with 30 ml of ethyl acetate. The organic layer was washed with 30 ml of brine, dried (MgSO_4), and then evaporated. The resulting residue was purified by silica gel column chromatography (NH-silica gel; n-hexane/ethyl acetate), to give 0.19 g (86%) of a free form of the title compound as a pale yellow oil.


$^1\text{H-NMR}$ (400Mz, CDCl_3) δ : 1.25-1.45 (4H, m), 1.52 (3H, t, $J = 7.2\text{Hz}$), 1.65-1.77 (2H, m), 1.87-1.95 (1H, m), 1.96-2.06 (2H, m), 2.65-2.73 (2H, m), 2.88-2.96 (2H, m), 3.22 (1H, dd, $J = 8\text{Hz}$, $J = 17.6\text{Hz}$), 3.54 (2H, s), 3.89 (3H, s), 4.18 (2H, q, $J = 7.2\text{Hz}$), 6.84 (1H, s), 7.16 (1H, s), 7.23-7.35 (5H, m).

ESI-MS: $m/z = 394$ ($\text{M}+\text{H}^+$)

Example 1

1-Benzyl-4-[(5,6-dimethoxy-2-chloro-1-indanon)-2-yl]methylpiperidine hydrochloride

[0043]



[0044] The following reaction was performed in nitrogen atmosphere

[0045] In 10 ml of tetrahydrofuran (THF) was dissolved 0.25 g (0.86 mmol) of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine obtained in accordance with Example 4 of JP-A 1-79151. After cooling to -78°C , 0.99 ml (0.96 mmol) of a 1.0 M-lithium bis(trimethylsilyl)amide/THF solution was added therein. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and 0.13 g of a (0.97 mmol) N-chlorosuccinimide/ THF (2 ml) solution and 0.5 ml of hexamethylphosphoramide (HMPA) were added thereto. After gradually elevating the temperature from -78°C to room temperature and stirring for 2 hours, an aqueous saturated ammonium solution (50 ml) was added and the mixture was extracted with ethyl acetate (50 ml). The organic layer was washed with 50 ml of brine, dried (MgSO_4), and then evaporated. The residue was purified by NH-silica gel column chromatography (n-hexane/ethyl acetate system) and then by preparative thin layer chromatography (methylene chloride/methanol system), to give 0.17 g (62%) of a free form of the title compound as a pale yellow oil. The oil was converted into a hydrochloride by a conventional method and recrystallized from ethanol/t-butyl methyl ether, to give the title compound as pale yellow crystals.

Melting point: 202 to 204°C .


$^1\text{H-NMR}$ (400Mz, CDCl_3) δ (ppm): 1.66-2.28 (7H, m), 2.60-2.74 (2H, m), 3.37-3.53 (4H, m), 3.92 (3H, s), 3.98 (3H, s), 4.13 (2H, dd, $J = 13.2\text{Hz}$, $J = 25.2\text{Hz}$), 6.80 (1H, s), 7.19 (1H, s), 7.42-7.47 (3H, m), 7.58-7.65 (2H, m), 12.42 (1H, bs).

ESI-MS: $m/z = 414$ ($\text{M}+\text{H}^+$).

Example 2

1-Benzyl-4-[(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methylpiperidine hydrochloride

[0046]



[0047] The following reaction was performed in nitrogen atmosphere.

[0048] In 10 ml of THF was dissolved 0.25 g (0.66 mmol) of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine. After cooling to -78°C , 0.99 ml (0.99 mmol) of a 1.0 M lithium bis(trimethylsilyl)amido/THF solution was added thereto. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and bromine (0.051 ml, 1.00 mmol)/THF (2 ml) solution was added thereto. After gradually elevating the temperature from -78°C to room temperature and stirring overnight, an aqueous saturated ammonium chloride solution (50 ml) was added and the mixture was extracted with ethyl acetate (50 ml). The organic layer was washed with brine (50 ml) dried (MgSO_4), and then evaporated. The residue was purified by NH-silica gel column chromatography (n-hexane/ethyl acetate system) and then by preparative thin layer chromatography (methylene chloride/methanol system), to give 0.071 g (24%) of a free form of the title compound as a pale yellow oil. The oil was converted into a hydrochloride by a conventional method and recrystallized from ethanol/n-butyl methyl ether, to give the titled compound as pale yellow crystals.

Melting point: 164 to 165°C .


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ (ppm) 1.74 (1H, dd, $J = 6.4\text{Hz}$, $J = 14.8\text{Hz}$), 1.89 (1H, d, $J = 13.2\text{Hz}$), 2.04-2.28(4H, m), 2.33 (1H, d, $J = 14.8\text{Hz}$), 2.65-2.80 (2H, m), 3.42-3.68 (4H, m), 3.92 (3H, s), 3.96 (3H, s), 4.13-4.26 (2H, m), 6.81 (1H, s), 7.22 (1H, s), 7.45 (3H, bs), 7.58-7.66 (2H, m), 12.05 (1H, bs).

ESI-MS; $m/z = 458$ ($\text{M}+\text{H}^+$).

Example 3

1-Benzyl-4-[(5,6-dimethoxy-2-hydroxy-1-indanon)-2-yl]methylpiperidine hydrochloride

[0049]



[0050] The following reaction was performed in nitrogen atmosphere.

[0051] In 10 ml of THF was dissolved (0.25 g, 0.66 mmol) of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine. After cooling to -78°C , 0.99 ml (0.99 mmol) of a 1.0 M lithium bis(trimethylsilyl)amido/THF solution was added thereto. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and 0.02 g (0.96 mmol) - (10-camphorsulfonyl)oxaziridin/THF (2 ml) solution was added thereto. After gradually elevating the temperature from -78°C to room temperature and stirring for 4 hours, an aqueous saturated ammonium chloride solution (50 ml) was added and the mixture was extracted with ethyl acetate (50 ml). The organic layer was washed with brine (50 ml), dried (MgSO_4), and then evaporated. The residue was purified by NH-silica gel column chromatography (n-hexane/ethyl acetate system), to give 0.26 g (quantitative amount) of a free form of the title compound as a pale yellow oil. The oil was converted into a hydrochloride by a conventional method and solidified by using diethyl ether, to give the title compound as a pale yellow amorphous.


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ (ppm) 1.56-2.20 (7H, m), 2.58-2.74 (2H, m), 3.06-3.50 (5H, m), 3.89 (3H, s), 3.97 (3H, s), 4.14 (2H, bs), 5.82 (1H, s), 7.11 (1H, s), 7.42 (3H, bs), 7.60 (2H, bs), 12.04 (1H, bs).

ESI-MS; $m/z = 396$ ($\text{M}+\text{H}^+$).

Example 4

Synthesis of 1-Benzyl-4-[(5,6-dimethoxy-2-methyl-1-indanon)-2-yl]methylpiperidine hydrochloride

[0052]



[0053] The following reaction was performed in nitrogen atmosphere.

[0054] In 10 ml of THF was dissolved 0.25 g (0.66 mmol) of 1-benzyl-4-[(5,6-dimethoxy-1-indanon)-2-yl]methylpiperidine. After cooling to -78°C , 0.99 ml (0.99 mmol) of a 1.0 M-lithium bis(trimethylsilyl)amide/THF solution was added thereto. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and 0.062 ml (0.99 mmol) of iodomethane was added thereto. After gradually elevating the temperature from -78°C to room temperature and stirring for 4 hours, an aqueous saturated ammonium chloride solution (50 ml) was added and the mixture was extracted with ethyl acetate (50 ml). The organic layer was washed with brine, dried (MgSO_4), and then evaporated. The residue was purified by preparative thin layer chromatography (methylene chloride/methanol system), to give 0.16 g (62%) of a free form of the title compound as a pale yellow oil. The oil was converted into a hydrochloride by a conventional method and recrystallized from ethanol/t-butyl methyl ether, to give the title compound as pale yellow crystals.

Melting point: 194 to 195°C .


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ (ppm) 1.19 (3H, s), 1.50-1.85 (5H, m), 2.00-2.14 (2H, m), 2.44-2.60 (2H, m), 2.63 (1H, d, J 17.2Hz), 2.99 (1H, d, J=17.2H), 3.35 (2H, bt, J=14H), 3.90 (3H, s), 3.97 (3H, s), 4.08 (2H, bs), 6.82 (1H, bs), 7.13 (1H, s), 7.35-7.50 (3H, m), 7.58 (2H, bs), 12.25 (1H, bs).

ESI-MS; $m/z = 394$ ($\text{M}+\text{H}^+$).

Example 5

Synthesis of 1-Benzyl-4-[(5,6-dimethoxy-2-azido-indanon)-2-yl]methylpiperidine hydrochloride

[0055]



[0056] In N,N-dimethylformamide (9 ml) was dissolved 0.30 g (0.65 mmol) of 1-benzyl-4-[(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methylpiperidine obtained in Example 2, followed by adding acetic acid (3 ml) and a solution of 0.21 g (3.23 mmol) of sodium azide/water (5 ml). After stirring at 70°C for 3 hours, the mixture was cooled down to room temperature and an aqueous saturated sodium carbonate solution (50 ml) was added. The resulting mixture was extracted with ethyl acetate (50 ml). The organic layer was washed with brine (50 ml), dried (MgSO_4), and then evaporated. The residue was purified by silica gel column chromatography (methylene chloride/methanol system), to give 0.18 g (65%) of a free form of the title compound as a pale yellow oil. The oil was converted into a hydrochloride by a conventional method and recrystallized from ethanol/t-butyl methyl ether, to give the title compound as pale yellow crystals. Melting point: 153 to 154°C .


$^1\text{H-NMR}$ (400MHz, CDCl_3): δ (ppm) 1.64-1.74 (1H, s), 1.82-1.94 (2H, m), 1.97-2.24 (4H, m), 2.64-2.78 (2H, m), 3.01 (1H, d, J = 17.2Hz), 3.19 (1H, d, J = 17.2H), 3.40-3.56 (2H, m), 3.91 (3H, s), 3.98 (3H, s), 4.20 (2H, dd, J = 13.2Hz, J = 19.6Hz), 6.82 (1H, s), 7.16 (1H, s), 7.40-7.47 (3H, m), 7.58-7.67 (2H, m).

ESI-MS; $m/z = 421$ ($\text{M}+\text{H}^+$).

Example 6

1-Benzyl-4-[(5,6-diethoxy-2-chloro-1-indanon)-2-yl]methylpiperidine hydrochloride

[0057]



This reaction was performed in nitrogen atmosphere. In 5 ml of THF was dissolved 0.10 g (0.26 mmol) of 1-benzyl-4-[(5,6-diethoxy-2-chloro-1-indanon)-2-yl]methylpiperidine. After cooling to -78°C , 0.38 ml (0.38 mmol) of a 1.0 M-lithium bis(trimethylsilyl)amide/THF solution was added thereto. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and 4 ml of THF solution containing 0.051 g (0.38 mmol) of N-chlorosuccinimide was added thereto. After gradually elevating the temperature from -78°C to room temperature and stirring for 4 hours, 30 ml of aqueous saturated ammonium chloride solution was added and the mixture was extracted with 30 ml of ethyl acetate. The organic layer was washed with 30 ml of brine, dried (MgSO_4), and then evaporated. The resulting residue was purified by preparative thin layer chromatography (methylene chloride/methanol), to give 0.058 g (51%) of a free form of the title compound as a pale yellow oil.


$^1\text{H-NMR}$ (400Mz, CDCl_3) δ : 1.34-1.48 (2H, m), 1.47 (3H, t, $J = 7.2\text{Hz}$), 1.51 (3H, t, $J = 7.2\text{Hz}$), 1.84-1.81 (4H, m), 1.90-2.03 (2H, m), 2.21 (1H, dd, $J = 4\text{Hz}$, $J = 14.4\text{Hz}$), 2.80-2.89 (2H, m), 3.45 (2H, s), 3.48 (2H, s), 4.11 (2H, q, $J = 7.2\text{Hz}$), 4.16 (2H, q, $J = 7.2\text{Hz}$), 6.79 (1H, s), 7.21 (1H, s), 7.22-7.33 (5H, m).

The free form was converted into hydrochloride and by a conventional method and recrystallized from ethanol/*t*-butyl methyl ether, to give the objective compound as pale yellow crystals. Melting point: 166 to 167°C . ESI-MS: $m/z = 442$ ($\text{M}+\text{H}^+$)

Example 7

1-Benzyl-4-[(5-ethoxy-6-methoxy-2-chloro-1-indanon)-2-yl]methylpiperidine hydrochloride

[0060]



This reaction was performed in nitrogen atmosphere. In 5 ml of THF was dissolved 0.14 g (0.35 mmol) of 1-benzyl-4-[(5-ethoxy-6-methoxy-1-indanon)-2-yl]methylpiperidine obtained in Reference Example 1. After cooling to -78°C , 0.52 ml (0.52 mmol) of a solution of 1.0 M-lithium bis(trimethylsilyl)amide/THF was added. After elevating the temperature from -78°C to -10°C over 30 minutes, the solution was again cooled down to -78°C and a solution of 0.070 g (0.52 mmol) of N-chlorosuccinimide in 6 ml of THF was added. After gradually elevating the temperature from -78°C to room temperature and stirring for 4 hours, 30 ml of an aqueous saturated ammonium chloride solution was added and the resulting mixture was extracted with ethyl acetate (30 ml). The organic layer was washed with 30 ml of brine, dried (MgSO_4), and then evaporated. The resulting residue was purified by preparative thin layer chromatography (methylene chloride/methanol), to give 0.077 g (52%) of a free form of the title compound as a pale yellow oil.

$^1\text{H-NMR}$ (400Mz, CDCl_3) δ : 1.35-1.50 (2H, m) 1.53 (3H, t, $J = 7.2\text{Hz}$), 1.64-1.82 (4H, m), 1.92-2.04 (2H, m), 2.21 (1H, dd, $J = 4\text{Hz}$, $J = 14.4\text{Hz}$), 2.82-2.90 (2H, m), 3.46 (2H, s), 3.49 (2H, s), 3.91 (3H, s), 4.19 (2H, q, $J = 7.2\text{Hz}$), 6.80 (1H, s), 7.23 (1H, s), 7.24-7.33 (5H, m).

The free form was converted into a hydrochloride by a conventional method and recrystallized from ethanol/*t*-butyl methyl ether, to give the objective title compound as pale yellow crystals. Melting point: 165 to 167°C .

ESI-MS: $m/z = 428$ ($\text{M}+\text{H}^+$)

Pharmacological Test Example

[0063] Hereinafter, pharmacological test examples for demonstrating usefulness of the compounds of the present invention as medicaments will be presented.

In vitro acetylcholinesterase inhibitory effect

[0064] Using rat brain homogenate as a source of acetylcholinesterase, esterase activity was assayed in accordance with the method of Ellman et al.¹⁾. That is, acetylthiocholine as a substrate, a test compound and DTNB (5,5'-dithiobis (2-nitrobenzoic acid)) were added to mouse brain homogenate, and after incubation, the produced thiocholine was reacted with DTNB to produce a yellow product. This was measured as a change in absorbance at 412 nm to obtain acetylcholinesterase activity. The acetylcholinesterase inhibiting activity of each test compound was obtained as 50% inhibition concentration (IC₅₀). Note that the test compounds were used after dissolving them in physiological saline.

¹⁾ Ellman G.L., Courtney, K.D., Andres, V. and Featherstone, R.M., (1961), *Biochem. Pharmacol.*, 7, 88-95.

Results


[0065]

Example	IC ₅₀ (nM)
1	2.0
6	2.0
Donepezil Hydrochloride	6.7

[0066] In the above test examples, the compounds of the present invention represented by the above formula (I), a salt thereof or hydrates thereof exhibited significant acetylcholinesterase inhibitory effect. The compounds (I) of the present invention are useful as an agent for treating, preventing or ameliorating various types of senile dementia, cerebrovascular dementia and attention deficit hyperactivity disorder. In consideration of activity, side effects, number of administration, form of administration etc. Especially, they are useful as an agent for treating, preventing or ameliorating Alzheimer-type senile dementia.


Claims


1. A compound represented by the formula:



a pharmacologically acceptable salt thereof or hydrates thereof.

In the formula, R¹ represents any group selected from the groups represented by the formulae:






and


(wherein R^3 's are the same as or different from each other and each represents hydrogen atom, a halogen atom, hydroxyl group, a C_{1-6} alkyl group, a C_{3-6} cycloalkyl group, a C_{1-6} alkoxy group, a C_{1-6} alkoxyalkoxy group, a halogeno C_{1-6} alkyl group, a hydroxy C_{1-6} alkyl group, a cyano C_{1-6} alkyl group, an amino C_{1-6} alkyl group, a halogeno C_{1-6} alkoxy group, a hydroxy C_{1-6} alkoxy group, a cyano C_{1-6} alkoxy group, a lower acyl group, nitro group, an optionally substituted amino group, an optionally substituted carbamoyl group, mercapto group or a C_{1-6} thioalkoxy group; R^4 represents hydrogen atom or a C_{1-6} alkyl group; R^5 represents a halogen atom (provided that fluorine is excluded), hydroxy group, a C_{1-6} alkyl group, a C_{1-6} alkoxy group, cyano group, a halogeno C_{1-6} alkyl group, a hydroxy C_{1-6} alkyl group, a cyano C_{1-6} alkyl group, an amino C_{1-6} alkyl group, nitro group, an azido group, an optionally substituted amino group, an optionally substituted carbamoyl group, an optionally substituted carboxyl group, mercapto group or a C_{1-6} thioalkoxy group; the partial structure --- represents a single bond or double bond; m is 0 or an integer from 1 to 6; n is an integer from 1 to 4; and p is an integer of 1 or 2); and

R^6 represents a C_{3-6} cycloalkyl/methyl, a 2,2-(alkylenedioxy)ethyl or a group represented by the formula.




wherein the ring A represents a benzene ring or a heterocyclic ring; (R^6)s are the same as or different from each other and each represents hydrogen, a halogen atom, hydroxyl group, nitrile group, a C_{1-6} alkyl group, a C_{3-8} cycloalkyl group, a C_{1-6} alkoxy group, a C_{1-6} alkoxyalkoxy group, an aryloxy group, an aralkyloxy group, a halogeno C_{1-6} alkyl group, a hydroxy C_{1-6} alkyl group, a cyano C_{1-6} alkyl group, a halogeno C_{1-6} alkoxy group, a hydroxy C_{1-6} alkoxy group, a cyano C_{1-6} alkoxy group, a lower acyl group, nitro group, an optionally substituted amino group, an optionally substituted amide group, mercapto group or a C_{1-6} thioalkoxy group, and two of the R^6 may together form an aliphatic ring, an aromatic ring, a heterocyclic ring or an alkylenedioxy ring; and q is 0 or an integer from 1 to 5.

2. The compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof, wherein R^1 is represented by the formula:



In which R^{3a} and R^{3b} are the same as or different from each other and each represents a C_{1-6} alkoxy group; and R^5 and m have the same meanings as defined above.

3. The compound according to Claim 2, a pharmacologically acceptable salt thereof or hydrates thereof, wherein R^{3a} and R^{3b} are methoxy groups.
4. The compound according to any of Claims 1 to 3, a pharmacologically acceptable salt thereof or hydrates thereof, wherein R^6 is chlorine or bromine.
5. The compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof, wherein m is 1.
6. The compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof, wherein R^2 is a group represented by the formula:



wherein the ring A, R^6 and q have the same meanings as defined above.

7. The compound according to Claim 6, a pharmacologically acceptable salt thereof or hydrates thereof, wherein the ring A is a benzene ring.
8. The compound according to Claim 6, a pharmacologically acceptable salt thereof or hydrates thereof, wherein the ring A is a pyridine ring.
9. The compound according to Claim 6, a pharmacologically acceptable salt thereof or hydrates thereof, wherein q

is an integer of 1 or 2.

10. The compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof which is the one selected from:

1-benzyl-4-[(5,6-dimethoxy-2-chloro-1-indanon)-1-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-bromo-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-iodo-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-hydroxy-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-methyl-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-ethyl-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-azido-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-amino-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-methylamino-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-dimethylamino-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-acetamido-1-indanon)-2-yl]methylpiperidine,
 1-benzyl-4-[(5,6-dimethoxy-2-methanesulfonamide-1-indanon)-2-yl]methylpiperidine,
 3-(1-benzylpiperidin-4-yl)-2-chloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone,
 3-(1-benzylpiperidin-4-yl)-2,2-dichloro-1-(2,3,4,5-tetrahydro-1H-1-benzoozepin-8-yl)-1-propanone,
 5,7-dihydro-3-(1-chloro-2-[(1-(phenylmethyl)-4-piperidinyl)ethyl]-6H-pyrido[4,5-f]-1,2-benzisooxazol-6-one,
 5,7-dihydro-3-(1,1-dichloro-2-[(1-(phenylmethyl)-4-piperidinyl)ethyl]-6H-pyrido[4,5-f]-1,2-benzisooxazol-6-one,
 1-(2-methyl-5-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2-chloro-1-propanone, and
 1-(2-methyl-6-benzothiazolyl)-3-[1-(phenylmethyl)-4-piperidinyl]-2,2-dichloro-1-propanone.

11. A medicament comprising the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof as an active element.

12. The medicament according to Claim 11, which is an acetylcholinesterase inhibitor.

13. A medicament according to Claim 11, which is an agent for treating, preventing or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder.

14. The medicament according to Claim 13, wherein the various types of senile dementia is Alzheimer type senile dementia.

15. A medical composition comprising a pharmacologically effective amount of the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof and a pharmacologically acceptable carrier.

16. A method of preventing, treating or ameliorating diseases against which the inhibition of an acetylcholinesterase is effective, by administering a pharmacologically effective amount of the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof to a patient.

17. A method of treating, preventing or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder, by administering a pharmacologically effective amount of the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof to a patient.

18. Use of the compound according to claim 1, a pharmacologically acceptable salt thereof or hydrates thereof for producing an agent for preventing, treating or ameliorating diseases against which the inhibition of an acetylcholinesterase is effective.

19. Use of the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof for producing an acetylcholinesterase inhibitor.

20. Use of the compound according to Claim 1, a pharmacologically acceptable salt thereof or hydrates thereof for producing an agent for preventing, treating or ameliorating various types of senile dementia, cerebrovascular dementia or attention deficit hyperactivity disorder.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP00/05968

A. CLASSIFICATION OF SUBJECT MATTER Int. Cl. ⁷ C07D211/32, 401/06, 498/04 103, 417/06, 413/06, A61K31/445, 55, 454, A61P25/28, 25/14, 43/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) Int. Cl. ⁷ C07D211/32, 401/06, 498/04 103, 417/06, 413/06, A61K31/445, 55, 454, A61P25/28, 25/14, 43/00		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) CA (STN), CAPLUS (STN), CAOLD (STN), REGISTRY (STN)		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
EX	WO, 00/51985, A1 (EISAI CO., LTD.), 08 September, 2000 (08.09.00) (Family: none) Full text	1-15, 18-20
X	JP, 5-268176, A (EISAI CO., LTD.), 14 October, 1997 (14.10.97) (Family: none) Full text; See especially the working example 5 on compounds	1-3, 5-7, 9, 11-15, 16-20 2-4, 8, 10
X	EP, 602242, A1 (YOSHITOMI PHARMACEUTICAL INDUSTRIES, LTD.), 22 June, 1994 (22.06.94), Full text; See especially the working example 48 on compounds	1, 5-7, 9, 11-15, 18-20 2-4, 8, 10
A	WO, 93/04663, A1 & JP, 5-320160, A & JP, 6-41125, A	
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubt on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed	"I" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, each combination being obvious to a person skilled in the art "Z" document member of the same patent family	
Date of the final completion of the international search 16 November, 2000 (16.11.00)		Date of mailing of the international search report 28 November, 2000 (28.11.00)
Name and mailing address of the ISA/ Japanese Patent Office Facsimile No.		Authorized officer Telephone No.

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP00/05968

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(c)(a) for the following reasons:

1. ☒ Claims Nos.: 16,17
because they relate to subject matter not required to be searched by this Authority, namely:
Claims 16 and 17 pertain to methods for treatment of the human body by therapy.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/210 (continuation of first sheet (1)) (July 1992)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/13427

C (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y, P	US 6,044,155 A (THOMLINSON et al.) 28 March 2000 (28.03.2000), column 4, lines 27-31, column 5, lines 23-29.	62

Form PCT/ISA/210 (continuation of second sheet) (July 1998)

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/13427

Continuation of B. FIELDS SEARCHED Item 3: EAST

search terms: random, pseudorandom, integrated circuit, chip, microchip, protected memory, audit, inspect, erase, reuse

Form PCT/ISA/210 (exam sheet) (July 1998)